



**Hospital onset COVID-19
mortality in Scotland**

7 March 2020 to 31 December 2020



24 February 2021

This is a management information publication

Published management information is non-official statistics which may be in the process of being transitioned into official statistics. They may not comply with the UK Statistics Authority's Code of Practice with regard to high data quality or high public value but there is a public interest or a specific interest by a specialist user group in accessing these statistics as there are no associated official statistics available.

Users should therefore be aware of the aspects of data quality and caveats surrounding these data, all of which are listed in this document.

Find out more about management information publications at:

<https://www.statisticsauthority.gov.uk/wp-content/uploads/2016/06/National-Statisticians-Guidance-Management-Information-and-Official-Statistics.pdf>

Contents

Introduction	4
Main points.....	5
Results and commentary	6
Implications for improved outcomes and infection prevention.....	11
References.....	11
Contact.....	12
Further information.....	12
Rate this publication.....	12
Appendices	13
Appendix 1 – Publication metadata	13
Appendix 2 – Early access details	19
Appendix 3 – Model results for adjusted hospital onset COVID-19 mortality (all-cause at 28 days).....	20

Introduction

Antimicrobial Resistance and Healthcare Associated Infection (ARHAI) Scotland, part of National Services Scotland, works closely with Public Health Scotland to deliver under the Health Protection Scotland (HPS) COVID-19 response. This second release provides data for COVID-19 hospital onset mortality in Scotland for the period 7 March 2020 to 31 December 2020. A report for period 1 July to 30 September was not published due to the small number of COVID-19 cases during this time.

Nosocomial transmission of SARS-CoV-2 contributes significantly to the overall burden of infection within these settings. Deaths occurring in patients with COVID-19 are an important measure of patient outcome. Therefore, monitoring COVID-19 mortality in hospital patients and publishing the data is critical in the development and monitoring of local and national improvement plans to improve patient outcomes, inform the development of infection prevention and control measures, shape policy and guide research. [Further information on the epidemiology of COVID-19 in healthcare settings can be found on the Health Protection Scotland website.](#)

A report on COVID-19 hospital onset cases is published weekly and is available from: <https://beta.isdscotland.org/find-publications-and-data/population-health/covid-19/hospital-onset-covid-19-cases-in-scotland/>

This report describes 28-day all-cause mortality in cases of COVID-19 that were identified during an inpatient stay in an NHS hospital in Scotland, including those cases which are thought to have developed the infection as a result of nosocomial transmission.

Main points

- Different methodologies are employed to measure mortality in cases of COVID-19. This report uses 28-day all-cause mortality in laboratory confirmed cases of COVID-19 rather than deaths where suspected or confirmed COVID-19 is listed on the death certificate. This is aligned with the daily reporting of deaths by Public Health Scotland.
- Overall more than a quarter of patients who were diagnosed with COVID-19 during an inpatient stay died within 28 days of their first positive test for SARS-CoV-2. Nearly a third of patients who developed probable or definite hospital onset COVID-19 died within 28 days.
- The difference in all-cause mortality between the onset categories can largely be explained by differences in the demographics of cases. Cases with probable or definite hospital onset COVID-19 are older than cases diagnosed in the first two days of admission to hospital and are likely to be sicker (although co-morbidity data was not available at the time of writing).
- After adjustment, patients who develop probable or definite hospital onset COVID-19 (day 8 of admission onwards) are at no greater risk of dying than inpatients who likely acquired SARS-CoV-2 in the with non-hospital onset (day 1 or 2 of admission).
- After adjustment for the confounding effects of hospital onset status, age and sex, patients who were first diagnosed with COVID-19 in hospital had lower odds of death within 28 days in wave 2 compared with wave 1.
- Older hospitalised patients are likely to have longer lengths of stay in hospital, increasing their risk of healthcare associated infection including COVID-19. These patients are also more likely to die from other causes and these are not distinguished in all-cause mortality estimates.
- Asymptomatic testing for SARS-CoV-2 has increased since the beginning of the pandemic. This will have increased case ascertainment including those with mild or asymptomatic disease who may have a lower risk of dying from COVID-19. It was not possible to distinguish between symptomatic cases and asymptomatic cases identified by testing policies nor control for this during these analyses.
- Preventing transmission of SARS-CoV-2 is critical to reducing morbidity and mortality from COVID-19, particularly in older hospital patients. Infection prevention and control precautions and early detection and management of cases is vital in efforts to reduce the spread of SARS-CoV-2 in hospital settings.

Results and commentary

COVID-19 deaths by hospital onset status

A total of 10,541 cases of COVID-19 were diagnosed during an inpatient stay in Scotland between the first case identified in hospitals on the 7 March and 31 December 2020 which includes 5,362 cases for the period July 2020 to December 2020. The total number of these patients who died within 28 days (all-cause) was 2,782 (26.4%) which includes 1,358 deaths in the period July 2020 to December 2020.

Cases of COVID-19 were categorised based on date of first positive SARS-CoV-2 PCR test following admission to a health board. Patients where the first positive sample was taken on day 1 or 2 of admission were likely to have acquired SARS-CoV-2 in the community. For patients where the first positive sample was taken on days 3-7 (indeterminate hospital onset), it is not possible to determine where acquisition was likely to have taken place e.g. in the community or during their hospital stay. Patients where the first positive samples were taken on days 8-14 (probable hospital onset) and days 15+ (definite hospital onset) are the categories where there is a likelihood of nosocomial transmission within the health board.

A total of 1,015 patients with probable or definite hospital onset COVID-19 died within 28 days of their first positive sample (29.9%). Mortality was highest among patients with probable (28.5%) or definite hospital onset COVID-19 (30.6%) ([Table 1](#)).

Table 1: Number of COVID-19 deaths by onset status in Scotland overall: specimen dates up to 31 December 2020.^{1,2,3}

Hospital onset status	Mar-Jun 2020 Mortality within 28 days (n)	Mar-Jun 2020 Mortality within 28 days (%)	Mar-Jun 2020 Total Cases	July-Dec 2020 Mortality within 28 days (n)	July-Dec 2020 Mortality within 28 days (%)	July-Dec 2020 Total Cases	Total Mortality within 28 days (n)	Total Mortality within 28 days (%)	Total Cases
Non-hospital onset (day 1 or 2 of admission)	919	26.1%	3,525	637	22.5%	2,833	1,556	24.5%	6,358
Indeterminate hospital onset (days 3-7)	73	24.2%	302	138	28.0%	492	211	26.6%	794
Probable hospital onset (days 8-14)	96	34.8%	276	189	26.1%	724	285	28.5%	1,000
Definite hospital onset (days 15+)	336	31.2%	1,076	394	30.0%	1,313	730	30.6%	2,389
Scotland	1,424	27.5%	5,179	1,358	25.3%	5,362	2,782	26.4%	10,541

1. Source of data is Electronic Communication of Surveillance in Scotland (ECOSS) data, the Rapid Admission Preliminary Inpatient Data (RAPID) data or local admission data, and National Records of Scotland (NRS).
2. The data used has not been adjusted for potential factors that may affect mortality e.g. severity of COVID-19 disease and patient comorbidities.
3. Cases diagnosed in the community (not during an inpatient stay) were excluded from these analyses

A logistic regression model was developed to adjust for confounding between the onset categories ([Appendix 3](#)). The model included all cases of COVID-19 identified during the hospital stay and the comparator group was selected as those cases thought to have acquired the infection in the community (day 1 or 2 of admission). After adjustment for the confounding effects of age, sex and pandemic wave, there was no significant difference between all-cause mortality in cases of indeterminate, probable and definite hospital onset COVID-19 compared with patients diagnosed with COVID-19 during the first 2 days of their admission to the health board (where the inpatient likely acquired the virus in the community). Age, sex and pandemic wave were all significantly associated with 28-day all-cause mortality. The model results are presented in [Appendix 3](#).

The pandemic wave was included as survival may have improved as knowledge of the course of the infection improved, or the severity of disease reduced over time. All-cause mortality decreased between wave 1 (specimen date $\leq 26/07/2020$) and wave 2 (specimen date $>26/07/2020$), after adjustment for case mix including the distribution of hospital onset cases. The reasons for this will be multifactorial and not intended to be explained by this model.

There are some limitations and caveats to these modelling analyses that must be considered in the interpretation. Age, sex and pandemic wave are currently the only risk factors with comprehensive data available and further modelling to adjust for other risk factors such as underlying co-morbidity and ethnicity will be undertaken as the data become available. Cases who are in the probable and definite category have a longer length of stay prior to developing COVID-19. This is indicative of underlying medical conditions which will be a risk factor for mortality and some of these patients may have died irrespective of COVID-19. Similarly, it is not known if some of the patients in the non-hospital onset category were emergency admissions due to a severe COVID-19 infection which has implications for the comparison with the probable and definite hospital onset mortality. Additionally, asymptomatic testing for SARS-CoV-2 has increased since the beginning of the pandemic. This will have increased case ascertainment including those with mild or asymptomatic disease who may have a lower risk of dying from COVID-19. This may affect comparisons in all-cause mortality across all onset groups and in particular the day 1 or 2 onset cases due to increased admission testing (though serial asymptomatic testing has been in place in some areas). It was not possible to distinguish between symptomatic cases and asymptomatic cases identified by testing policies and control for this during these analyses. There are some early indications of increased disease severity in people infected with variant of concern (VOC) 202012/01 (also known as B.1.1.7) compared to people with non-VOC virus variants.¹ If there is further definitive evidence of increased mortality, the impact of the increasing incidence of VOC202012/01 may require to be considered in future mortality analyses.

The distribution of 28-day all-cause mortality by age and sex for each of the hospital onset categories is described in [Table 2](#). The highest all-cause mortality was reported in male patients and in older age groups, where risk of death increased with increasing age. This is

in line with the outputs from the logistic regression model. These data are unadjusted for underlying co-morbidity and other risk factors and therefore should be interpreted with due caution. The median age of patients who died following a probable or definite hospital onset COVID-19 diagnosis (79 years) was significantly higher than those patients with likely community acquisition (day 1 or 2 of admission) (71 years, unadjusted $p < 0.001$).

Information relating to the specialty or setting where these cases are being cared for is not currently available but will be included in future analyses as these data become available. This will be essential for developing local and national focused improvement plans.

Table 2: COVID-19 deaths within 28 days, by onset status, age group and sex: specimen dates up to 31 December 2020. ^{1,2,3}

Age Group / Hospital onset status	Female mortality (n)	Female cases (n)	Female mortality (%)	Male mortality (n)	Male cases (n)	Male mortality (%)	Total mortality (n)	Total cases (n)	Total mortality (%)
0-24	1	123	0.8%	0	114	0.0%	1	237	0.4%
Non-Hospital Onset	1	109	0.9%	0	94	0.0%	1	203	0.5%
Indeterminate Hospital Onset	0	5	0.0%	0	8	0.0%	0	13	0.0%
Probable Hospital Onset	0	1	0.0%	0	2	0.0%	0	3	0.0%
Definite Hospital Onset	0	8	0.0%	0	10	0.0%	0	18	0.0%
25-44	9	293	3.1%	11	291	3.8%	20	584	3.4%
Non-Hospital Onset	5	231	2.2%	6	228	2.6%	11	459	2.4%
Indeterminate Hospital Onset	1	21	4.8%	2	22	9.1%	3	43	7.0%
Probable Hospital Onset	1	15	6.7%	1	9	11.1%	2	24	8.3%
Definite Hospital Onset	2	26	7.7%	2	32	6.3%	4	58	6.9%
45-64	110	984	11.2%	186	1,313	14.2%	296	2,297	12.9%
Non-Hospital Onset	67	726	9.2%	125	989	12.6%	192	1,715	11.2%
Indeterminate Hospital Onset	9	71	12.7%	10	73	13.7%	19	144	13.2%
Probable Hospital Onset	13	60	21.7%	17	90	18.9%	30	150	20.0%
Definite Hospital Onset	21	127	16.5%	34	161	21.1%	55	288	19.1%
65-74	200	868	23.0%	360	1,279	28.1%	560	2,147	26.1%
Non-Hospital Onset	121	534	22.7%	240	817	29.4%	361	1,351	26.7%
Indeterminate Hospital Onset	11	53	20.8%	24	96	25.0%	35	149	23.5%
Probable Hospital Onset	16	86	18.6%	28	107	26.2%	44	193	22.8%
Definite Hospital Onset	52	195	26.7%	68	259	26.3%	120	454	26.4%
75-84	396	1,426	27.8%	615	1,579	38.9%	1,011	3,005	33.6%
Non-Hospital Onset	210	738	28.5%	353	886	39.8%	563	1,624	34.7%
Indeterminate Hospital Onset	29	108	26.9%	42	123	34.1%	71	231	30.7%
Probable Hospital Onset	38	167	22.8%	58	168	34.5%	96	335	28.7%
Definite Hospital Onset	119	413	28.8%	162	402	40.3%	281	815	34.5%
85+	422	1,295	32.6%	472	976	48.4%	894	2,271	39.4%
Non-Hospital Onset	199	551	36.1%	229	455	50.3%	428	1,006	42.5%
Indeterminate Hospital Onset	45	131	34.4%	38	83	45.8%	83	214	38.8%
Probable Hospital Onset	57	172	33.1%	56	123	45.5%	113	295	38.3%
Definite Hospital Onset	121	441	27.4%	149	315	47.3%	270	756	35.7%
Grand Total	1,138	4,989	22.8%	1,644	5,552	29.6%	2,782	10,541	26.4%

1. Source of data is Electronic Communication of Surveillance in Scotland (ECOSS) data, the Rapid Admission Preliminary Inpatient Data (RAPID) data or local admission data, and National Records of Scotland (NRS).
2. The data used has not been adjusted for potential factors that may affect mortality e.g. severity of COVID-19 disease and patient comorbidities.
3. Cases diagnosed in the community (not during an inpatient stay) were excluded from these analyses

Comparison with other mortality data in Scotland

Nearly a third of cases of hospital onset COVID-19 (probable and definite) died within 28 days of the first positive specimen (29.9%). Cases who are in the probable and definite category have, by the design of the case definition, a longer length of stay prior to developing COVID-19. This is indicative of underlying medical condition which will also be a risk factor for mortality and some of these patients may have died irrespective of COVID-19. It is not possible to quantify this with the data currently available and comparison with other published mortality data is difficult due to the different ways in which the deaths are defined and the populations in which they are reported.

All-cause mortality data is available for other infection types commonly associated with nosocomial infection. In 2018, nearly a fifth of cases (18.2%) of *Staphylococcus aureus* bacteraemia; 10.9% of *Clostridioides difficile* cases aged 15 years and older; and 13.7% of cases of *Escherichia coli* bacteraemia had died within 30 days of their diagnosis.² These mortality estimates are not directly comparable as the population of cases will include hospitalised and non-hospitalised cases and the duration of follow up differs.

The COVID-19 analyses are restricted to hospitalised patients and it would be anticipated that these patients are sicker and require hospital care. Crude mortality rates in Scotland, used in the calculation of hospital standardised mortality ratios, for January to September 2020 indicate that 4.7% and 10.5% of patients aged 60-79 years and 80+ years, respectively die within 30 days of an admission to hospital.³ A UK study of mortality in nosocomial COVID-19 in older people reported that 27.0% of cases of nosocomial COVID-19 (diagnosed 15 or more days after admission) had died within seven days.⁴ After adjustment, nosocomial mortality was reduced compared with community acquired COVID-19 in hospitalised patients.

Any comparisons between differing mortality measures must be treated with caution. All-cause mortality includes deaths where COVID-19 may not have been either the underlying or contributory cause of death. In addition, deaths due to COVID-19 infection of long duration will be underestimated in 28-day all-cause mortality e.g. cases who have died more than 28 days after their first positive sample would not be included using 28-day all-cause mortality. All-cause mortality is not subject to the same biases as death certification that are introduced as a result of subjectivity or changes in the way deaths were registered during the early stages of the pandemic. In addition, death certification includes cases of presumed COVID-19 without a positive SARS-CoV-2 test and these are not included in 28-day all-cause mortality measure.

Implications for improved outcomes and infection prevention

As SARS-CoV-2 is a new and emerging pathogen, new international evidence around effective treatments, diagnostics, and prevention and control are being published at a rapid pace. Within Scotland and across the UK there are a number of organisations reviewing the evidence, contributing to the research and ensuring that key measures are reflected in the COVID-19 national response guidance.

As 28-day all-cause mortality is no different in patients admitted to hospital with likely community acquisition to those who develop probable or definite hospital onset COVID-19, preventing transmission of SARS-CoV-2 in all settings is critical to reducing morbidity and mortality from COVID-19.

A continued focus on the broader public health interventions along with the application of infection prevention and control precautions in line with current guidance will reduce the risk of transmission. In healthcare settings, early detection and appropriate management through testing of symptomatic patients and screening of asymptomatic patients with early identification of any contacts, particularly when there is an unexpected case or outbreak, is vital in efforts to reduce the spread of SARS-CoV-2 in these settings.

References

- (1) NERVTAG update. 2021 [cited 2021 Feb 17]; Available from: URL: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/961042/S1095_NERVTAG_update_note_on_B.1.1.7_severity_2_0210211.pdf
- (2) Antimicrobial Resistance and Healthcare Associated Infection Scotland. Healthcare Associated Infections. 2019 Annual report. ARHAI Scotland 2020 [cited 2021 Feb 9]; Available from: URL: <https://www.hps.scot.nhs.uk/web-resources-container/healthcare-associated-infection-annual-report-2019/>
- (3) Public Health Scotland. Hospital Standardised Mortality Ratios. PHS 2021 February 9 [cited 2021 Feb 9]; Available from: URL: <https://beta.isdscotland.org/find-publications-and-data/health-services/hospital-care/hospital-standardised-mortality-ratios/>
- (4) Carter B, Collins JT, Barlow-Pay F, Rickard F, Bruce E, Verduri A, et al. Nosocomial COVID-19 infection: examining the risk of mortality. The COPE-Nosocomial Study (COVID in Older PEople). J Hosp Infect 2020 Oct;106(2):376-84.

Contact

Laura Imrie

Clinical Lead, Antimicrobial Resistance and Healthcare Associated Infections (ARHAI) Scotland

NHS National Services Scotland

Phone: 0141 300 1922

Email: nss.hpshaic@nhs.scot

Further information

Further Information can be found on the [HPS website](#).

For more information on types of infections included in this report, please see the [COVID-19](#) pages on the HPS website.

The next release of this publication will be 26 May 2021 (subject to additional cases in the intervening period).

Rate this publication

Please [provide feedback](#) on this publication to help us improve our services.

Appendices

Appendix 1 – Publication metadata

Metadata indicator	Description
Publication title	Hospital onset COVID-19 mortality in Scotland
Description	This release provides information on hospital onset COVID-19 mortality, there is a need for consistent reporting using standardised case definitions.
Theme	Infections in Scotland
Topic	COVID-19
Format	Word document
Data source(s)	<p>COVID-19 Cases:</p> <p>Case data source: Electronic Communication of Surveillance in Scotland (ECOSS)</p> <p>Admissions data Source: Rapid Admission Preliminary Inpatient Data (RAPID) or Local Patient Admissions Systems</p> <p>Mortality data source: National Records of Scotland (NRS)</p>
Date that data are acquired	2 February 2021
Release date	24 February 2021
Frequency	Quarterly (subject to additional cases in the intervening period).
Timeframe of data and timeliness	<p>Timeframe of this publication was decided by first positive sample in Scotland which fell into anyone of the hospital onset categories (i.e. 7 March 2020)</p> <p>The latest iteration of data is 31 December 2020, therefore the data are 7 or 8 weeks in arrears.</p> <p>No report was produced in November 2020 to allow sufficient data to accumulate for information governance needs and analysis to be done since the period of the last published report (March-June 2020).</p>
Continuity of data	Quarterly updates (subject to additional cases in the intervening period).
Revisions statement	These data are not subject to planned major revisions. However, ARHAI Scotland aims to continually improve the interpretation of the data and therefore analysis methods are regularly reviewed and may be updated in the future.

<p>Revisions relevant to this publication</p>	<p>Hospital onset data are continually validated by NHS boards, NRS and within the ECOSS laboratory database. Any changes to cases taken within hospital settings, which are validated by NHS boards, are tracked by ARHAI Scotland. Changes to retrospective mortality data for the hospital onset groups are outlined in the table below if applicable.</p> <table border="1" data-bbox="419 416 1412 902"> <thead> <tr> <th data-bbox="419 416 751 562">Hospital onset mortality as previously reported (26 August 2020)</th> <th data-bbox="751 416 1078 562">Hospital onset mortality as currently reported</th> <th data-bbox="1078 416 1412 562">Reason</th> </tr> </thead> <tbody> <tr> <td data-bbox="419 562 751 902">332 definite hospital onset cases with mortality identified for time period March-June 2020</td> <td data-bbox="751 562 1078 902">336 definite hospital onset cases with mortality identified for time period March-June 2020</td> <td data-bbox="1078 562 1412 902">4 community cases reassigned to definite hospital onset died within 28 days of first COVID-19 result. Details of all changes to hospital onset categories can be found in the weekly Hospital Onset COVID-19 Report.</td> </tr> </tbody> </table>	Hospital onset mortality as previously reported (26 August 2020)	Hospital onset mortality as currently reported	Reason	332 definite hospital onset cases with mortality identified for time period March-June 2020	336 definite hospital onset cases with mortality identified for time period March-June 2020	4 community cases reassigned to definite hospital onset died within 28 days of first COVID-19 result. Details of all changes to hospital onset categories can be found in the weekly Hospital Onset COVID-19 Report.
Hospital onset mortality as previously reported (26 August 2020)	Hospital onset mortality as currently reported	Reason					
332 definite hospital onset cases with mortality identified for time period March-June 2020	336 definite hospital onset cases with mortality identified for time period March-June 2020	4 community cases reassigned to definite hospital onset died within 28 days of first COVID-19 result. Details of all changes to hospital onset categories can be found in the weekly Hospital Onset COVID-19 Report.					
<p>Concepts and definitions</p>	<p>The transmission of COVID-19 is thought to occur mainly through respiratory droplets and through contact with contaminated surfaces. Further information on the epidemiology of COVID-19 in healthcare settings can be found on the Health Protection Scotland website. As sustained community transmission has occurred as the pandemic has progressed, it has become more challenging to identify true cases of hospital transmission.</p> <p>A system for monitoring COVID-19 is critical to tracking nosocomial transmission in healthcare settings to inform infection, prevention and control measures.</p> <p>Deaths occurring in patients with COVID-19 are an important measure of patient outcome. Therefore, monitoring COVID-19 mortality in hospital patients and publishing the data is critical to improve care of patients, inform the development of infection prevention and control measures, shape policy and guide research.</p> <p>The data provided are national data for Scotland representing the 14 NHS boards and one NHS special health board.</p> <p>The agreed nosocomial case definition for the UK is based on the number of days since admission to an NHS health board to the date of specimen sampling for a positive SARS-CoV-2 RT-PCR test. Time since admission to specimen sampling is categorised as:</p> <ul style="list-style-type: none"> • community onset (first positive specimen taken in the community), non-hospital onset (first positive specimen on day 1 or 2 of admission to NHS board); • indeterminate (first positive specimen on days 3 to 7 of admission to NHS board); • probable (first positive specimen on days 8 to 14 of admission to NHS board); and • definite hospital onset (first positive specimen date was 15 or more days after admission to NHS board). 						

Note that for the purposes of this report, cases diagnosed in the community (not during an inpatient stay) were excluded from these analyses to restrict the comparisons within the hospitalised patient population.

These definitions are necessary due to the maximum incubation period of 14 days for COVID-19 (see table below):

Day of sampling post admission	Nosocomial categorisation
Before admission	Community onset COVID-19 (not included in this report)
Day 1 of admission/on admission to NHS board	Non-hospital onset COVID-19
Day 2 of admission	Non-hospital onset COVID-19
Day 3 of admission	Indeterminate hospital onset COVID-19
Day 4 of admission	Indeterminate hospital onset COVID-19
Day 5 of admission	Indeterminate hospital onset COVID-19
Day 6 of admission	Indeterminate hospital onset COVID-19
Day 7 of admission	Indeterminate hospital onset COVID-19
Day 8 of admission	Probable hospital onset COVID-19
Day 9 of admission	Probable hospital onset COVID-19
Day 10 of admission	Probable hospital onset COVID-19
Day 11 of admission	Probable hospital onset COVID-19
Day 12 of admission	Probable hospital onset COVID-19
Day 13 of admission	Probable hospital onset COVID-19
Day 14 of admission	Probable hospital onset COVID-19
Day 15 of admission and onwards to discharge	Definite hospital onset COVID-19
Post discharge	Community onset COVID-19 (not included in this report)

The hospital onset cases in this report represent cases presenting in hospital and do not include COVID-19 associated with hospital care that present on readmission to hospital or post-discharge.

Start point of duration

Admission to health board was agreed as the appropriate point to start counting the duration of hospital stay to first positive specimen date, rather than the date of admission to a single hospital, since patients can be transferred between hospitals which would lead to restarting the clock to 'day 1' each time and therefore underestimating the number of nosocomial infections.

	<p>Any discharges and re-admissions which occur within the same calendar day will be classed as a continuous stay; the clock will not be restarted in these instances, only when a readmission occurs on the second day or more after any discharge.</p> <p>For definite, probable, indeterminate and non-hospital onset (day 1 or 2 of in-patient stay), the NHS board reported is where the first sample was taken, established either using Rapid Admission Preliminary Inpatient Data (RAPID) data and validated by the boards, or using individual NHS board's internal admissions systems. Since the definition of hospital-onset COVID-19 was determined using date of admission to NHS board, the board assigned may not represent the board of attribution of hospital-onset COVID-19 infection (Table above).</p> <p>Minimum data required for hospital onset COVID-19 cases to be validated:</p> <ul style="list-style-type: none"> - CHI number - Date of positive SARS-CoV-2 RT-PCR test - Date of admission to health board when patient tested positive for COVID-19 - NHS board where first positive test undertaken <p><i>Mortality definition</i></p> <p>In this report, all-cause mortality within 28 days of the COVID-19 diagnosis (laboratory specimen date) is used. Therefore, the data includes deaths where COVID-19 may not have been either the underlying or contributory cause of death. All-cause mortality depends solely on the number of deaths identified, and is not subject to bias that may be introduced as a result of inaccuracies in completion of the death certificate or coding of the cause of death. Using 28-days as the time period makes the assumption that most deaths related to COVID-19 will occur within this timeframe. Deaths occurring after this time period are more difficult to assess as being specifically related to COVID-19, though they are known to occur. Therefore, care should be taken when interpreting this data and when comparing published data on COVID-19 mortality that use different definitions.</p> <p><i>Wave 1 / Wave 2 definition</i></p> <p>The 26th of July was chosen as an arbitrary cut-off signifying the end of Wave 1 in Scotland, since cases of Hospital onset COVID-19 were at their lowest point over the course of the pandemic. Data from the 27th July onwards are considered Wave 2.</p>
Relevance and key uses of the statistics	Surveillance data are essential for monitoring trends and assisting in outbreak investigations and to understand the extent of ongoing transmission within the hospital setting. HPS offers support to NHS boards across Scotland to aid their local COVID-19 prevention strategies.
Accuracy	It is acknowledged that patients can be transferred between NHS health boards and if transferred into a different health board during the same hospital stay, then the clock would be restarted to 'day 1' which could lead to an underestimation of cases. However, the decision to restrict start date to admission to a single NHS health board represents the requirement to report at the health board-level. Any discharges and re-admissions which occur within the same calendar day will be overlooked - the clock will not be restarted in these instances, only when a readmission occurs on the second or more day after any discharge.

	<p>COVID-19 cases identified after discharge from hospital but within 14 days may be associated with the hospital. These cases, including those identified on readmission to hospital, are not included as hospital onset. This may result in under-reporting of COVID-19 cases associated with hospital care.</p> <p>All-cause mortality depends solely on the number of deaths identified, and is not subject to bias that may be introduced as a result of inaccuracies in completion of the death certificate or coding of the cause of death.</p>
Completeness	<p>Surveillance data are collected using the ECOSS system that allows data collectors in NHS boards to validate ECOSS records as well as identifying additional cases that may not be included in the Electronic Communications of Surveillance in Scotland (ECOSS) system. This therefore means that completeness is near to 100%. For mortality data, sufficient time is allowed for all cases to be followed up for 28 days. Some delays in reporting of death may occur but this will be minimal and completeness is near to 100%.</p>
Comparability	<p>The agreed nosocomial case definition for the UK has been adopted to allow comparison across the four nations. However, geographical differences for example NHS board versus NHS Trust have to be considered.</p> <p>The case numbers presented here are only for those COVID-19 cases who are positive as inpatients with an admission to an NHS hospital, i.e. community cases are excluded. Additionally, the end dates are different to that of the weekly report (i.e. week end 3rd Jan vs. 31st December). The data are therefore not wholly comparable with the weekly reporting of hospital onset cases.</p> <p>https://beta.isdscotland.org/find-publications-and-data/population-health/covid-19/hospital-onset-covid-19-cases-in-scotland/</p> <p>Deaths within this report are taken from the same data source used by Public Health Scotland (PHS) in their daily reported deaths, therefore comparisons are possible between these two sources. However, it should be noted that this report includes data until 31st December 2020 using COVID-19 specimen date, with deaths followed-up until 27th January (28 days later), whereas Publish Health Scotland report using date of death. Therefore, comparisons between each surveillance system should be treated with caution.</p>
Accessibility	<p>It is the policy of HPS to make its web sites and products accessible according to published guidelines.</p>
Coherence and clarity	<p>Tables and charts are accessible via the HPS website at: https://www.hps.scot.nhs.uk/a-to-z-of-topics/covid-19/hospital-onset-covid-19-cases-in-scotland/</p>
Value type and unit of measurement	<p>At national level, the number and proportion of COVID-19 cases who died within 28-days (all-cause) of a COVID-19 diagnosis are classed as definite hospital onset, probable hospital onset, indeterminate hospital onset, and non-hospital onset. The data are further broken down by age group and sex. For adjusting the data to account for confounding, pandemic wave is also included in combination with age group, sex and hospital onset status.</p>
Disclosure	<p>The HPS protocol on Statistical Disclosure Control Protocol is followed.</p>
Official Statistics designation	<p>Management Information</p>
UK Statistics Authority Assessment	<p>Not assessed</p>
Last published	<p>26 August 2020</p>

Next published	26 May 2021 (subject to additional cases in the intervening period)
Date of first publication	26 August 2020
Help email	nss.hpshaic@nhs.scot
Date form completed	24 February 2021

Appendix 2 – Early access details

Pre-Release Access

Under terms of the "Pre-Release Access to Official Statistics (Scotland) Order 2008", ARHAI Scotland is obliged to publish information on those receiving Pre-Release Access ("Pre-Release Access" refers to statistics in their final form prior to publication). The standard maximum Pre-Release Access is five working days. Shown below are details of those receiving standard Pre-Release Access.

Standard Pre-Release Access:

Scottish Government Health Department

NHS board Chief Executives

NHS board Communication leads

Appendix 3 – Model results for adjusted hospital onset COVID-19 mortality (all-cause at 28 days)

Univariable and multivariable logistic regression results on 28-day mortality outcome of COVID-19 cases.¹

Variable		Alive within 28 days (%)	Mortality within 28 days (%)	OR ² (univariable)	OR ² (multivariable)
Sex	F	3851 (77.2)	1138 (22.8)	-	-
	M	3908 (70.4)	1644 (29.6)	1.42 (1.30-1.55, p<0.001)	1.59 (1.45-1.74, p<0.001)
Age group (years)	0-49	1098 (95.4)	53 (4.6)	-	-
	50-59	1069 (89.1)	131 (10.9)	2.54 (1.84-3.56, p<0.001)	2.44 (1.76-3.42, p<0.001)
	60-69	1275 (79.3)	332 (20.7)	5.39 (4.03-7.36, p<0.001)	5.18 (3.87-7.08, p<0.001)
	70-79	1913 (70.4)	805 (29.6)	8.72 (6.60-11.75, p<0.001)	8.66 (6.55-11.69, p<0.001)
	80+	2404 (62.2)	1461 (37.8)	12.59 (9.58-16.91, p<0.001)	13.14 (9.98-17.68, p<0.001)
Hospital onset status	Non-hospital onset	4802 (75.5)	1556 (24.5)	-	-
	Indeterminate hospital onset	583 (73.4)	211 (26.6)	1.12 (0.94-1.32, p=0.196)	0.97 (0.81-1.16, p=0.744)
	Probable hospital onset	715 (71.5)	285 (28.5)	1.23 (1.06-1.43, p=0.006)	0.97 (0.83-1.14, p=0.750)
	Definite hospital onset	1659 (69.4)	730 (30.6)	1.36 (1.22-1.51, p<0.001)	1.02 (0.91-1.14, p=0.704)
Pandemic wave	Wave 1 (≤26/07/20)	3772 (72.6)	1424 (27.4)	-	-
	Wave 2 (>26/07/20)	3987 (74.6)	1358 (25.4)	0.90 (0.83-0.98, p=0.020)	0.84 (0.77-0.92, p<0.001)

¹ Note that for the purposes of this report, cases diagnosed in the community (not during an inpatient stay) were excluded from these analyses to restrict the comparisons within the hospitalised patient population.

² OR = Odds ratio